

Integral cohomology of classical groups over a finite field

Denis Jeandupeux

Department of Mathematics, Cornell University, White Hall, Ithaca, NY 14853, USA

Communicated by A. Dold

Received 14 September 1991

Revised 30 March 1992

Abstract

Jeandupeux, D., Integral cohomology of classical groups over a finite field, *Journal of Pure and Applied Algebra* 84 (1993) 43–58.

Let G be one of the groups GL_n , SL_n , U_n , SU_n or Sp_{2n} over a finite field of characteristic p . We calculate the order of the integral Chern classes, obtained by Brauer lifting. From the cellular cochains of suitable Eilenberg–Mac Lane spaces, we construct a complex. Using a description of the mod l cohomology of G (l prime $\neq p$), we prove that the homology of this complex is the integral cohomology of G , away the p -torsion.

Quillen gives the mod l cohomology of GL_n and Fiedorowicz and Priddy give this cohomology for the groups U_n and Sp_{2n} . We compute this cohomology for the special groups SL_n and SU_n .

Introduction

Let \mathbb{F}_q be a finite field with q elements of characteristic p . The purpose of this paper is to describe the integral cohomology, away the p -torsion, of the following groups of matrices (we refer to [5] for the definitions and further details):

the general linear group $GL_n = GL_n(\mathbb{F}_q)$ and the special linear group $SL_n = SL_n(\mathbb{F}_q)$ over \mathbb{F}_q ,

the unitary group $U_n = U_n(\mathbb{F}_{q^2})$ and the special unitary group $SU_n = SU_n(\mathbb{F}_{q^2})$ over \mathbb{F}_{q^2} and

the symplectic group $Sp_{2n} = Sp_{2n}(\mathbb{F}_q)$ ($\subset GL_{2n}$).

Let G denote one of these groups and BG be the associated classifying space. For each integral Chern class $\tilde{c}_i \in H^{2i}(BG; \mathbb{Z})$, obtained by Brauer lifting, we calculate its order s_i and then we choose a subcomplex C_i of the cellular cochain of the Eilenberg–Mac Lane space $K(\mathbb{Z}/s_i, 2i-1)$ such that H^*C_i is the poly-

Correspondence to: Professor D. Jeandupeux, Résidence de la Côte 40, CH-1110 Morges, Switzerland.

mial subalgebra $\mathbb{Z}[\tilde{x}_i]/(s_i\tilde{x}_i)$ of $H^*(K(\mathbb{Z}/s_i, 2i-1); \mathbb{Z})$, where \tilde{x}_i is a generator of $H^{2i}(K(\mathbb{Z}/s_i, 2i-1); \mathbb{Z}) \simeq \mathbb{Z}/s_i$. Our main result is the following:

Theorem. $\tilde{H}^*(\bigotimes_{i \in I} C_i)$ is isomorphic to $\tilde{H}^*(BG; \mathbb{Z}[\frac{1}{p}])$ as abelian groups, where I depends on G .

In [9], Huebschmann gives a description of that cohomology in the stable case, i.e. for $G_\infty = \varinjlim G_n$.

Our proof uses a description of the mod l cohomology of G , for each prime l distinct from p , with generators the Chern classes and other classes e_i , $i \in \mathbb{N}$, considered first by Quillen in [13]. The first part of the paper gives this description. For GL_n this is done in [13]. For U_n and Sp_{2n} we use the complete description of this cohomology given by Fiedorowicz and Priddy in [5].

For SL_n we have two cases. If l divides the order of GL_1 we study the multiplicative structure of the Lyndon–Hochschild–Serre spectral sequence associated to the exact sequence $1 \rightarrow SL_n \rightarrow GL_n \xrightarrow{\det} GL_1 \rightarrow 1$. In particular, we show that the spectral sequence collapses and the GL_1 -action on $H^*(SL_n; \mathbb{Z}/l)$ is trivial. If l does not divide the order of GL_1 we use some basic tools of cohomology of groups and we adapt the detection theorems of Quillen (cf. [12]) to our situation.

It turns out that the same method works for SU_n .

The orthogonal groups are not treated here. This case seems to be somewhat different from the others, especially about the 2-torsion (cf. [5, I.7]). Moreover, the method used to compute the cohomology of the symplectic groups does not work. Indeed the homomorphism induced by the inclusion $O_2(\mathbb{F}_q) \rightarrow GL_2(\mathbb{F}_q)$ in mod 2 cohomology is not surjective (cf. Theorem 1.3 and [5, IV.4 and IV.8]). On the other hand there are different kinds of orthogonal groups and it would be better to treat them in another paper.

1. Mod l cohomology

Let k be an integer and r the smallest integer ≥ 1 such that $k^r \equiv 1 \pmod{l}$. For any space or group X we note $H^*X = H^*(X; \mathbb{Z}/l)$. By $P[x_1, x_2, \dots]$ (resp. $\Lambda[x_1, x_2, \dots]$) we mean the polynomial ring (resp. exterior algebra) over \mathbb{Z}/l with the generators x_1, x_2, \dots . We consider the classical fibration

$$JU(k) \xrightarrow{\phi} BU \xrightarrow{1-\psi^k} BU, \quad (1)$$

where $1-\psi^k$ is the application representing the Adams operation in K -theory. We know that $H^*BU = P[c'_1, c'_2, \dots]$ where $c'_i \in H^{2i}BU$ is the i th universal Chern class. We put $c_i = \phi^*c'_i \in H^{2i}JU(k)$. Quillen shows in [13] that $c_i = 0$ for $i \not\equiv 0 \pmod{r}$ and for $i \equiv 0 \pmod{r}$ he constructs classes $e_i \in H^{2i-1}JU(k)$. The case

where $l \neq 2$ or if $l = 2$ then $k \equiv 1 \pmod{4}$ is called the typical one and the case where $l = 2$ and $k \equiv 3 \pmod{4}$ is the exceptional one.

Theorem 1.1 (Quillen). *The monomials*

$$c_r^{\alpha_1} c_{2r}^{\alpha_2} \cdots e_r^{\beta_1} e_{2r}^{\beta_2} \cdots$$

with $\alpha_i \in \mathbb{N}$ and $\beta_i \in \{0, 1\}$ form a basis for $H^*JU(k)$. In the typical case $(e_{jr})^2 = 0$ and there is an algebra isomorphism

$$H^*JU(k) \cong P[c_r, c_{2r}, \dots] \otimes \Lambda[e_r, e_{2r}, \dots].$$

In the exceptional case,

$$e_j^2 = \sum_{0 \leq i < j} c_i c_{2j-1-i}. \quad \square$$

Let G_n denote GL_n or U_n . Let $k = q$ for $G_n = GL_n(\mathbb{F}_q)$ and $k = -q$ for $G_n = U_n(\mathbb{F}_{q^2})$. Let G_∞ be the direct limit $\varinjlim G_n$. The space $JU(k)$ has the homotopy type of the space BG_∞^+ obtained by performing the ‘+’ construction on the classifying space BG_n (cf. [5, 7, 13]). We consider the canonical map

$$\varepsilon : BG_n \rightarrow BG_\infty \rightarrow BG_\infty^+ \sim JU(k).$$

We write also c_i and e_i for the image of these classes under ε^* or the homomorphisms induced by canonical inclusions of groups.

Theorem 1.2. *The algebra homomorphism $\varepsilon^* : H^*JU(k) \rightarrow H^*BG_n$ is surjective. The kernel is the ideal generated by e_{jr} and c_j for $jr > n$. In particular, the monomials*

$$c_r^{\alpha_1} c_{2r}^{\alpha_2} \cdots c_{mr}^{\alpha_m} e_r^{\beta_1} e_{2r}^{\beta_2} \cdots e_{mr}^{\beta_m},$$

with $\alpha_i \in \mathbb{N}$, $\beta_i \in \{0, 1\}$ and $m = [n/r]$, form a basis of H^*G_n .

Let SG_n denote the groups SL_n or SU_n .

Theorem 1.3. *The algebra homomorphism $H^*G_n \rightarrow H^*SG_n$, induced by the inclusion, is surjective. The kernel is the ideal generated by e_1 and c_1 . For $r \geq 2$ it is an isomorphism.*

Let d be the smallest integer such that $q^{2d} \equiv 1 \pmod{l}$.

Theorem 1.4. *The algebra homomorphism $H^*GL_{2n} \rightarrow H^*Sp_{2n}$, induced by the*

inclusion, is surjective. If $r = 2d$, it is an isomorphism. If $r = d$, the kernel is the ideal generated by $e_{(2j+1)r}$ and $c_{(2j+1)r}$, for $j \in \mathbb{N}$.

In both cases, we have

$$H^* \mathrm{Sp}_{2n} = P[c_{2d}, c_{4d}, \dots, c_{2dm}] \otimes \Lambda[e_{2d}, e_{4d}, \dots, e_{2dm}]$$

for $1 \leq n \leq \infty$, where $m = \lfloor n/2d \rfloor$.

2. Integral cohomology away the p -torsion

Let $\tilde{c}_i' \in H^{2i}(BU; \mathbb{Z})$ be the i th integral universal Chern class. We consider the fibration (1) and the canonical map $\varepsilon : BG_n \rightarrow BG_\infty^+ \sim JU(k)$. We write \tilde{c}_i for the image of \tilde{c}_i' under the homomorphisms Φ^* , $\varepsilon^* \circ \Phi^*$ and those induced by inclusions of groups.

Theorem 2.1. *We have:*

- (1) For G_n and $1 \leq i \leq n$, the order of \tilde{c}_i is $|k^i - 1|$.
- (2) For SG_n and $2 \leq i \leq n$, the order of \tilde{c}_i is $|k^i - 1|$ and \tilde{c}_1 is zero.
- (3) For $i > n$, $\tilde{c}_i = 0$.

Let G denote G_n or SG_n and let I be the set of the i 's such that $1 \leq i \leq n$ for G_n and $2 \leq i \leq n$ for SG_n . For each $i \in I$, let s_i denote the order of \tilde{c}_i in $H^{2i}(BG; \mathbb{Z})$.

We consider the canonical fibration of Eilenberg–Mac Lane spaces

$$K(\mathbb{Z}/s_i, 2i-1) \xrightarrow{\beta} K(\mathbb{Z}, 2i) \xrightarrow{s_i} K(\mathbb{Z}, 2i), \quad (2)$$

where s_i is induced by the multiplication by s_i on \mathbb{Z} .

We denote by the same letter a cohomology class and its corresponding map under the usual isomorphism $[X, K(R, m)] \simeq H^m(X; R)$.

In [13], Quillen builds a class $\tilde{e}_i \in H^{2i}(BG; \mathbb{Z}/s_i)$ such that

$$\beta \tilde{e}_i = \tilde{c}_i, \quad (3)$$

where β denote the Bockstein homomorphism associated to the coefficient exact sequence

$$0 \rightarrow \mathbb{Z} \xrightarrow{s_i} \mathbb{Z} \rightarrow \mathbb{Z}/s_i \rightarrow 0.$$

Thus

$$\gamma_i = \tilde{e}_i : BG \rightarrow K(\mathbb{Z}/s_i, 2i-1)$$

is a lifting of \tilde{c}_i . We put $K_i = K(\mathbb{Z}/s_i, 2i-1)$.

The generator \tilde{x}_i of $H^{2i}(K_i; \mathbb{Z}) \simeq \mathbb{Z}/s_i$ generates a polynomial subalgebra $P_i = \mathbb{Z}[\tilde{x}_i]/(s_i\tilde{x}_i)$ of $H^*(K_i; \mathbb{Z})$ and P_i is a direct summand (cf. [4]).

Lemma 2.2. *We can choose \tilde{x}_i such that $\gamma_i^* \tilde{x}_i = \tilde{c}_i$.*

Proof. Let $\iota \in H^{2i}(K(\mathbb{Z}, 2i); \mathbb{Z})$ be the fundamental class. The Serre cohomology sequence of (2) shows that $\beta^* \iota$ is a generator of $H^{2i}(K_i; \mathbb{Z})$. Then by (3), $\gamma_i^*(\beta^* \iota) = \beta \gamma_i = \beta \tilde{c}_i = \tilde{c}_i$. \square

We denote by $(C(X), d)$ the cellular cochain complex of a space X . We chose a cocycle $z_i \in C^{2i}(K_i)$ which represents \tilde{x}_i and $b_i \in C^{2i-1}(K_i)$ such that $db_i = s_i z_i$. Let C_i be the subcomplex of $C(K_i)$ given by

$$C_i^{2im} = \mathbb{Z} z_i^m, \quad C_i^{2im-1} = \mathbb{Z} z_i^{m-1} \cup b_i, \quad C_i^j = 0 \text{ otherwise,}$$

where \cup is the cup product of the cochains.

Remark. The homomorphism $H^* C_i \rightarrow H^*(K_i; \mathbb{Z})$, induced by the inclusion, is injective with image P_i .

Let $g : BG \rightarrow \prod_{i \in I} K_i$ be the map given by the γ_i 's and

$$h : \bigotimes_{i \in I} C_i \rightarrow \bigotimes_{i \in I} C(K_i) \simeq C\left(\prod_{i \in I} K_i\right) \xrightarrow{g^*} C(BG)$$

be the homomorphism of complexes obtained by the Eilenberg–Zilber equivalence.

Theorem 2.3. *The homomorphism*

$$h^* : \tilde{H}^*\left(\bigotimes_{i \in I} C_i\right) \rightarrow \tilde{H}^*(BG; \mathbb{Z}[\frac{1}{p}])$$

is an isomorphism of abelian groups.

We consider the canonical homomorphism

$$f : \bigotimes_{i=1}^n C_{2i} \rightarrow \bigotimes_{i=1}^{2n} C_i \xrightarrow{h} C(BGL_{2n}) \rightarrow C(BSp_{2n}).$$

Theorem 2.4. *The homomorphism*

$$f^* : \tilde{H}^*\left(\bigotimes_{i=1}^n C_{2i}\right) \rightarrow \tilde{H}^*(BSp_{2n}; \mathbb{Z}[\frac{1}{p}])$$

is an isomorphism of abelian groups.

Corollary 2.5. *The order of the integral Chern class $\tilde{c}_{2i} \in H^{2i}(BSp_{2n}; \mathbb{Z})$ is $q^{2i} - 1$ for $1 \leq i \leq n$. The other classes are zero. \square*

Remarks. (1) The theorems remain true in the stable case, i.e. for $n = \infty$. So they describe $\tilde{H}^*(BG_\infty; \mathbb{Z})$, $\tilde{H}^*(BSG_\infty; \mathbb{Z})$ and $\tilde{H}^*(BSp_\infty; \mathbb{Z})$ since these groups have no p -torsion (cf. [7, 13] and (11)).

(2) Let R and R' be the subalgebras generated by the x_i 's and \tilde{c}_i 's respectively. Then $h^* : R \rightarrow R'$ is an isomorphism of algebras.

(3) The groups $H^k(\bigotimes_{i \in I} C_i)$ are easily computed by the Künneth formula.

Example. Let $G = SL_4(\mathbb{F}_5)$. The mod l cohomology is not trivial only for $l = 2, 3, 13$ or 31 ($l \neq 5$). The involved Chern classes are \tilde{c}_2, \tilde{c}_3 and \tilde{c}_4 , their orders are 24, 124 and 624 respectively. Then we have $H^{4j}C_2 = \mathbb{Z}/24z_2^j$, $H^{6j}C_3 = \mathbb{Z}/124z_3^j$, $H^{8j}C_4 = \mathbb{Z}/624z_4^j$ and the other groups are zero. By applying the Künneth formula to compute $H^*(C_2 \otimes C_3)$ and $H^*(C_2 \otimes C_3 \otimes C_4)$ successively, we obtain the integral cohomology groups of G (away the 5-torsion):

$$H^4G = \mathbb{Z}/24, \quad H^6G = \mathbb{Z}/124, \quad H^8G = \mathbb{Z}/24 \otimes \mathbb{Z}/624,$$

$$H^9G = \mathbb{Z}/4, \quad H^{10}G = \mathbb{Z}/4, \quad H^{11}G = \mathbb{Z}/4,$$

$$H^{12}G = \mathbb{Z}/4 \otimes \mathbb{Z}/24 \otimes \mathbb{Z}/124, \quad \dots$$

The nonlisted groups are zero.

3. Proof of Theorem 1.2

In [13], Quillen proves the theorem for $G = GL_n$. For $G = U_n$ we use the results of [5] and [13]. If $n < r$, l does not divide the order of U_n and then the mod l cohomology of U_n is trivial. For $n = r$ the l -Sylow subgroup C of U_r is cyclic and its order is l^a with $a = v_l((-q)^r - 1)$. We get a commutative diagram

$$\begin{array}{ccccc} BC & \xrightarrow{i} & BU_r & \longrightarrow & JU(-q) \\ B_f \downarrow & & & & \downarrow \phi \\ \prod_{i=1}^r BC & \xrightarrow{B_g} & \prod_{i=1}^r BU(1) & \longrightarrow & BU(r) \longrightarrow BU \end{array}$$

where i is induced by inclusion, $f(x) = (x, x^\alpha, x^{\alpha^2}, \dots, x^{\alpha^{r-1}})$ with $\alpha = q$ if r is even and $\alpha = q^2$ if r is odd, and $g = \prod_{i=1}^r \rho$ where $\rho : C \rightarrow S^1 = U(1)$ is the canonical representation (cf. [5]). Now we compute $\varepsilon^* i^* c_{jr}$ and $\varepsilon^* i^* e_{jr}$ as in the proof of Proposition 1 in [13]. In mod l cohomology i^* is injective and the Poincaré series of H^*BU_r is $(1 + t^{2r-1})/(1 - t^{2r})$. The result follows from the multiplicative structure of H^*BC .

For $n > r$ the map $BU_x \rightarrow JU(-q)$ induces an isomorphism in mod l cohomology. By [5] the canonical homomorphism $(U_r)^m \xrightarrow{\oplus} U_n$ (resp. $U_n \rightarrow U_x$) induces a monomorphism (resp. an epimorphism) in mod l cohomology. The proof for $n = r$ and the same computation as in the proof of Lemma 8 in [13], using the product formula, imply the theorem. \square

4. Proof of Theorem 1.3 in the case $r = 1$

The exact sequence

$$1 \rightarrow \mathrm{SG}_n \rightarrow G_n \xrightarrow{\det} C \rightarrow 1 \quad (4)$$

splits and G_n is the semi-direct product $\mathrm{SG}_n \rtimes C$, is the cyclic group G_1 . Let Y be the usual free resolution of \mathbb{Z} over $\mathbb{Z}C$ and let X be the standard resolution of \mathbb{Z} over $\mathbb{Z}\mathrm{SG}_n$ (cf. [3, I.5]). By a result of [11], $X \otimes Y$ is a free resolution of \mathbb{Z} over $\mathbb{Z}G_n$ and the cohomology of G_n is the homology of the bigraded differential algebra $B = \mathrm{Hom}_C(Y, \mathrm{Hom}_{\mathrm{SG}_n}(X; \mathbb{Z}/l))$. There is a filtration on B defined by $F^r(B^n) = \bigoplus_{k=r}^n B_{k, n-k}$ where $B_{p,q} = \mathrm{Hom}_C(Y_p, \mathrm{Hom}_{\mathrm{SG}_n}(X_q; \mathbb{Z}/l))$ and $B^n = \bigoplus_{p+q=n} B_{p,q}$. The corresponding spectral sequence E is the Lyndon–Hochschild–Serre spectral sequence of (4) (cf. [3, VII] and [11]). For $x \in H^*G_n$ we note \bar{x} the image of x in $E_x \simeq \mathrm{Gr}H^*G_n$.

Proposition 4.1. *For this spectral sequence we have:*

- (1) *The filtration degree of the monomial $c_1^{\alpha_1} c_2^{\alpha_2} \cdots c_n^{\alpha_n} e_1^{\beta_1} e_2^{\beta_2} \cdots e_n^{\beta_n}$ is $2\alpha_1 + \beta_1$ and the monomials $\bar{c}_1^{\alpha_1} \bar{c}_2^{\alpha_2} \cdots \bar{c}_n^{\alpha_n} \bar{e}_1^{\beta_1} \bar{e}_2^{\beta_2} \cdots \bar{e}_n^{\beta_n}$ of degree $p + q = 2(\alpha_1 + \cdots + \alpha_n) + \beta_1 + \cdots + \beta_n$ with $p = 2\alpha_1 + \beta_1$ form a basis of $E_x^{p,q}$.*
- (2) *$E_x^{p,q} = E_2^{p,q}$ for all p and q .*
- (3) *The C -action on $H^*\mathrm{SG}_n$, induced by the C -action on SG_n , is trivial.*

Now, since $E_2^{p,q} \simeq H^p(C; H^q\mathrm{SG}_n)$, we have $E_x^{0,q} \simeq H^q\mathrm{SG}_n$ and Theorem 1.3 follows from Proposition 4.1.

Proof of Proposition 4.1. We proceed by induction on q . Since the sequence (4) splits, $\det^*: H^*C \rightarrow H^*G_n$ is injective and then $\mathrm{Im} \det^* = E_x^{*,0} \subset H^*G_n$ is isomorphic to H^*G_1 . Now the C -action on $H^0\mathrm{SG}_n$ is trivial and $E_x^{*,0}$ is a quotient of $E_2^{*,0} \simeq H^*(G_1; H^0\mathrm{SG}_n) = H^*G_1$. This proves the proposition for $q = 0$. Suppose the proposition is true for $s \leq q - 1$ and write $b(m)$ for a monomial of the form $c_2^{\alpha_2} \cdots c_n^{\alpha_n} e_2^{\beta_2} \cdots e_n^{\beta_n}$ in H^mG_n . From $H^qG_n \simeq \bigoplus_{r+s=q} E_x^{r,s}$ and the multiplicative structure of E_x we deduce:

$$\begin{aligned} &\text{The filtration degree of the monomials } c_1^{\alpha_1} e_1^{\beta_1} b(q) \text{ is } 2\alpha_1 + \beta_1 \\ &\text{and their images } \bar{c}_1^{\alpha_1} \bar{e}_1^{\beta_1} \bar{b}(q) \text{ in } E_x^{2\alpha_1 + \beta_1, q} \text{ are linearly independent; moreover, the } \bar{b}(q) \text{ form a basis of } E_x^{0,q}. \\ &\text{In particular, the multiplication by } \bar{c}_1^{\alpha_1} \bar{e}_1^{\beta_1} \text{ is injective in } E_x. \end{aligned} \quad (5)$$

The differentials $d_r^{p,q}$ are zero for $r \geq 2$. In particular,

$$E_x^{0,q} = E_2^{0,q} \quad \text{and} \quad E_x^{1,q} = E_2^{1,q}. \quad (6)$$

Lemma 4.2. *The C -action on H^qSG_n is trivial.*

Hence by the universal coefficient theorem we have $E_2^{p,q} = E_2^{p,0} \otimes E_2^{0,q}$. From (5) and (6) it follows that $E_x^{p,q} = E_2^{p,q}$ for all p . This proves Proposition 4.1. \square

Proof of Lemma 4.2. We have an isomorphism of algebras

$$E_1^{*,q} \simeq \text{Hom}_C(Y_*; H^qSG_n). \quad (7)$$

Since $Y_p = \mathbb{Z}C$ for all p , we get a commutative diagram.

$$\begin{array}{ccccc} E_1^{0,q} & \xrightarrow{d_1^{0,q}} & E_1^{1,q} & \xrightarrow{d_1^{1,q}} & E_1^{2,q} \\ \parallel & & \parallel & & \parallel \\ H^qSG_n & \xrightarrow{g-1} & H^qSG_n & \xrightarrow{N} & H^qSG_n \end{array} \quad (8)$$

where g is a generator of C and $N = 1 + g + g^2 + \cdots + g^t$ with $t = |C| - 1$.

Claim 4.3. *The multiplication $E_1^{0,q} \otimes E_1^{1,0} \rightarrow E_1^{1,q}$ is an isomorphism.*

Proof. By an argument of dimension it suffices to prove the surjectivity. Since $E_1^{1,0} = E_2^{1,0}$ we can consider the class $e_1 \in E_2^{1,0} = H^1C$ as the cocycle in $E_1^{1,0}$ defined by $e_1(1) = 1 \in H^0SG_n = \mathbb{Z}/I$. Using the usual diagonal approximation associated with the resolution Y (cf. [3, V.1]) we get

$$(-1)^q(\varphi \cup e_1) = \varphi \quad \text{for } \varphi \in \text{Hom}_C(Y; H^qSG_n). \quad (9)$$

The claim follows now from (7). \square

Thus we can write every element in $E_1^{1,q}$ as $\varphi \cdot e_1$ for an unique $\varphi \in E_1^{0,q}$. Now if $x \in H^qSG_n$ is not fixed under the C -action then there exists a $\varphi \in E_1^{0,q}$ such that

$$d_1^{0,q}x = \varphi \cdot e_1 \neq 0 \quad (10)$$

and then $d_1^{1,q}d_1^{0,q}x = (d_1^{0,q}\varphi) \cdot e_1 = 0$. From (8) and (9) one sees that φ is fixed in H^qSG_n . Since $(H^qSG_n)^C \simeq E_2^{0,q} \subset E_1^{0,q}$ and $E_x^{0,q} = E_2^{0,q}$ by (6), $\bar{\varphi}$ is non-zero in E_x . Since \bar{e}_1 is also non-zero in E_x , $\bar{\varphi} \cdot \bar{e}_1$ is non-zero by (5). This contradicts (10) and proves Lemma 4.2. \square

Remark. Instead of using the complex B we can obtain a similar proof by taking

the cellular cochain complex of $BC = K(C, 1)$ with value in the system of local coefficients $H^q SG_n$.

5. Proof of Theorem 1.3 (continued)

In this section we prove Theorem 1.3 for $SG_n = SL_n$ if $r > 1$ and for $SG_n = SU_n$ if r is odd > 1 . The idea is to follow the method of [13, Corollary of Theorem 3].

Since $r > 1$, l is odd. Let t be q for SL_n and q^2 for SU_n . The l -Sylow subgroup C of SG_r is cyclic of order l^a with $a = \nu_l(t^r - 1)$ and is given as follows.

For SL_r we consider \mathbb{F}_{q^r} as an r -dimensional vector space over \mathbb{F}_q and we get a monomorphism $h : \mathbb{F}_{q^r}^* \rightarrow GL_r(\mathbb{F}_q)$ given by $h(a)x = ax$.

For SU_r we consider the standard Hermitian space $\mathbb{F}_{q^{2r}}$ (with form H) as an r -dimensional Hermitian space over \mathbb{F}_{q^2} with the form $\text{tr} \circ H$, where $\text{tr} : \mathbb{F}_{q^{2r}} \rightarrow \mathbb{F}_{q^2}$ is the trace map (cf. [5], note that r odd is required). As above we get a monomorphism $h : U_1(\mathbb{F}_{q^{2r}}) \rightarrow U_r(\mathbb{F}_{q^2})$ given by $h(a)x = ax$.

Then C is the p -primary component of $F_{q^r}^*$ for SL_n , resp. $U_1(\mathbb{F}_{q^{2r}})$ for SU_n , and $h(C)$ is contained in SG_r since $r > 1$.

We put $n = mr + e$ with $0 \leq e < r$ and we obtain a monomorphism

$$i : C^m \rightarrow SG_n, \quad x = (x_1, \dots, x_m) \mapsto (h(x_1), \dots, h(x_m), I_e).$$

Proposition 5.1. *The homomorphism $i^* : H^* SG_n \rightarrow H^* C^m$ is injective.*

The Galois group $\pi = \text{Gal}(\mathbb{F}_{q^r}/\mathbb{F}_q)$ is cyclic of order r generated by the Frobenius homomorphism $F : x \mapsto x^q$ and it acts on C . We get then an action of the wreath product $\pi^m \rtimes \Sigma_m$ on C^m , where the symmetric group Σ_m permutes canonically the factors.

Proposition 5.2. *In SG_n the action of any element of $\pi^m \rtimes \Sigma_m$ on C^m is realized by an inner automorphism.*

An inner automorphism induces the identity in cohomology and thus we have a monomorphism

$$i^* : H^* SG_n \rightarrow (H^* C^m)^{\pi^m \rtimes \Sigma_m}.$$

Now we consider the canonical homomorphism.

$$j^* : H^* G_n \rightarrow H^* SG_n \xrightarrow{i^*} (H^* C^m)^{\pi^m \rtimes \Sigma_m}$$

and Theorem 1.3 follows from the following proposition.

Proposition 5.3. *The homomorphism $j^* : H^*G_n \rightarrow (H^*C^m)^{\pi^m \rtimes \Sigma_m}$ is an isomorphism.*

Proof of Proposition 5.1. Since the index of SG_{mr} in SG_n is relatively prime to l , it suffices to show that C^m detects the mod l cohomology of SU_{mr} .

We consider the embedding $\Sigma_m \rightarrow G_{mr}$ such that Σ_m acts by conjugation on the diagonal matrices of m blocks $r \times r$ by permuting these blocks in the obvious way. The index of the subgroup $H_{m,r} = \Sigma_m \cap SG_{mr}$ of Σ_m ($\subset G_{mr}$) is 1 or 2 and, since l is odd, the wreath product $(SG_r)^m \rtimes H_{m,r}$ detects the cohomology of SG_{mr} . But the l -Sylow subgroup C of SG_r detects the cohomology of SG_r , and the following lemma implies Proposition 5.1. \square

Lemma 5.4. *Let G be a subgroup of SG_r . Suppose the mod l cohomology of G is detected by a subgroup K of G . Then the mod l cohomology of the wreath product $G^m \rtimes H_{m,r}$ ($\subset SG_{mr}$) is detected by K^m .*

Proof. By induction on m . There is nothing to prove for $m = 1$. We have two cases.

(1) If m is relatively prime to l then the index of $(G^{m-1} \rtimes H_{m-1,r}) \times G$ in $G^m \rtimes H_{m,r}$ is relatively prime to l and thus this subgroup detects the cohomology of $G^m \rtimes H_{m,r}$. The lemma follows by induction and the Künneth formula.

(2) If l divides m , we write $m = \bar{m}l$ and we consider the subgroup

$$(G^l \rtimes H_{l,r})^{\bar{m}} \rtimes H_{\bar{m},lr} \cong G^m \rtimes (H_{l,r}^{\bar{m}} \rtimes H_{\bar{m},lr})$$

of $G^m \rtimes H_{m,r}$. By induction its cohomology is detected by $(K^l)^{\bar{m}} = K^m$. But its index is relatively prime to l and so this subgroup detects the cohomology of $G^m \rtimes H_{m,r}$. This proves Lemma 5.4. \square

Proof of Proposition 5.2. We have $\Sigma_m \subset G_{mr} \subset G_n$. Moreover, $F \in G_r$ and for each $x \in C$ we have $F \circ h(x) \circ F^{-1} = h(F(x))$. So, in G_n , $\pi^m \rtimes \Sigma_m$ acts on C^m by inner automorphisms. Now for each $x \in G_n$ there exists b in $\mathbb{F}_{q^r}^*$ (or $U_1(\mathbb{F}_{q^{2r}})$) such that $\det h(b) = \det x^{-1}$, since $\det \circ h$ is surjective. But $\mathbb{F}_{q^r}^*$ and $U_1(\mathbb{F}_{q^{2r}})$ are cyclic and then $x \cdot i(b, 1, \dots, 1) \in SG_n$ acts as x on C^m . \square

Proof of Proposition 5.3. Quillen proved it for GL_n in [13]. For U_n , we remark that $j : C^m \rightarrow SU_n \rightarrow U_n$ factorizes by

$$C^m \rightarrow (U_r)^m \rightarrow U_n.$$

The proof of Theorem 1.2 (or [13, Section 5]) shows that $H^*U_r \rightarrow (H^*C)^\pi$ and $H^*U_n \rightarrow ((H^*U_r)^{\otimes m})^{\Sigma_m}$ are isomorphisms. \square

6. Proof of Theorem 1.4

Let $i : \mathrm{Sp}_{2n} \rightarrow \mathrm{GL}_{2n}$ be the inclusion. We write $2n = rs + e = 2dm + v$ with $0 \leq e < r$ and $0 \leq v < 2d$. For $r = 2d$, l does not divide the index of Sp_{2n} in GL_{2n} and then i^* is injective. We have $m = s$ and by comparing the Poincaré series of $H^*\mathrm{GL}_{2n}$ and $H^*\mathrm{Sp}_{2n}$ we see that i^* is surjective. For $r = d$, we first show that the theorem is true for $n = d$.

Lemma 6.1. $H^*\mathrm{Sp}_{2n} = P[c_{2d}] \otimes \Lambda[e_{2d}]$.

Then we consider the canonical map

$$f : (\mathrm{Sp}_{2d})^m \xrightarrow{j_1} \mathrm{Sp}_{2dm} \xrightarrow{j_2} \mathrm{Sp}_{2n} \rightarrow \mathrm{GL}_{2n} \rightarrow \mathrm{GL}_\infty.$$

By Lemma 6.1 we can write $H^*(\mathrm{Sp}_{2d})^m = P[x_1, \dots, x_m] \otimes \Lambda[y_1, \dots, y_m]$, where x_i (resp. y_i) corresponds to the class c_{2d} (resp. e_{2d}) of the i th factor. Using the notation of Lemma 9 in [13] and the product formula we show by induction on m that for each integer k

$$f^*c_{2dk} = \sigma_k, \quad f^*e_{2dk} = d\sigma_k, \quad f^*c_{(2k+1)d} = f^*e_{(2k+1)d} = 0.$$

In particular, $f^*(e_{2dk})^2 = 0$ (cf. [13, p. 565]). By Theorem 1.2 the image of $H^*\mathrm{GL}_{2n}$ in $H^*(\mathrm{Sp}_{2d})^m$ is then $P[\sigma_1, \dots, \sigma_m] \otimes \Lambda[d\sigma_1, \dots, d\sigma_m]$ and the theorem follows from the fact that j_1^* and j_2^* are monomorphisms (cf. [5]). \square

Proof of Lemma 6.1. For $d = 1$, $\mathrm{Sp}_2 = \mathrm{SL}_2$ and the lemma follows from Theorem 1.3 and the fact that $e_2^2 = 0$. If $d > 1$ then l is odd and we consider

$$f : \mathrm{GL}_d \rightarrow \mathrm{Sp}_{2d} \rightarrow \mathrm{GL}_{2d}$$

defined by

$$\mathrm{GL}_d \xrightarrow{\Delta} \mathrm{GL}_d \times \mathrm{GL}_d \xrightarrow{1 \times g} \mathrm{GL}_d \times \mathrm{GL}_d \xrightarrow{h} \mathrm{GL}_{2d} \xrightarrow{c} \mathrm{GL}_{2d},$$

where Δ is the diagonal map, $g(x) = (x^{-1})^t$, $h(x, y) = \mathrm{diag}(x, y)$ and c is the conjugation sending the $2d \times 2d$ matrix $\begin{pmatrix} 0 & I_d \\ -I_d & 0 \end{pmatrix}$ to $\mathrm{diag}(A, \dots, A)$ where $A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \in \mathrm{SL}_2$. Now c^* is the identity and $H^*\mathrm{GL}_d = P[c_d] \otimes \Lambda[e_d]$ since $r = d$ and l is odd.

Lemma 6.2. $f^*c_{2d} = \delta(c_d)^2$ and $f^*e_{2d} = \delta c_d e_d$ with $(\delta, q^d - 1) = 1$.

Hence f^* maps the subalgebra $P[c_{2d}] \otimes \Lambda[e_{2d}]$ of $H^*\mathrm{GL}_{2d}$ injectively in $H^*\mathrm{GL}_d$ and this subalgebra is isomorphic to $H^*\mathrm{Sp}_{2d}$ by [5]. This proves Lemma 6.1. \square

Proof of Lemma 6.2. The integral Chern class \tilde{c}_d generates a direct summand $\mathbb{Z}/q^d - 1$ in the group $H^{2d}(\mathrm{GL}_d; \mathbb{Z})$ (cf. [10] or Theorem 2.3). Hence $g^*\tilde{c}_d = \delta\tilde{c}_d$ with $(\delta, q^d - 1) = 1$. Since $H^{2d-1}(\mathrm{GL}_d; \mathbb{Z})$ is a p -group (cf. same ref.), the Bockstein homomorphism $\beta : H^{2d-1}(\mathrm{GL}_d; \mathbb{Z}/l) \rightarrow H^{2d}(\mathrm{GL}_d; \mathbb{Z})$ is injective and the same relation holds for e_d . The lemma follows from the product formula (cf. [13, p. 563]). \square

7. Proof of Theorem 1.3 (end)

It remains to prove the case r even for $\mathrm{SG}_n = \mathrm{SU}_n$. We have $r = 2d$ (cf. Section 1 for the notation) and l odd. If $n < r$, l does not divide the order of SU_n and then the mod l cohomology of SU_n is trivial.

For $n = r$ by [5] we have the following monomorphisms:

$$\mathrm{Sp}_{mr}(\mathbb{F}_q) \xrightarrow{g} \mathrm{SU}_{mr}(\mathbb{F}_{q^2}) \xrightarrow{h} U_{mr}(\mathbb{F}_{q^2}).$$

Since the indices of these subgroups are relatively prime to l , g^* and h^* are injective. By comparing the Poincaré series we see that $g^* \circ h^*$ is then an isomorphism, so g^* and h^* are isomorphisms.

For $n = mr + e$, $0 \leq e < r$, we consider the commutative diagram of inclusions

$$\begin{array}{ccc} \mathrm{SU}_{mr} & \xrightarrow{f} & \mathrm{SU}_n \\ h \downarrow & & \downarrow g \\ U_{mr} & \xrightarrow{k} & U_n \end{array}$$

We know that h^* and k^* are isomorphisms and by an argument of transfer f^* and g^* are injective. Hence f^* and g^* are isomorphisms. \square

8. Proof of Theorem 2.1

We first prove the following result:

Proposition 8.1. *The class $\tilde{c}_i = \Phi^* \tilde{c}'_i$ has order $|k^i - 1|$ in $H^{2i}(\mathrm{JU}(k); \mathbb{Z})$. In particular, \tilde{c}_i has no p -torsion.*

Proof. Since $\Omega \mathrm{BU}$ has the same homotopy type as U , we get the fibration

$$(*) \quad U \rightarrow \mathrm{JU}(k) \xrightarrow{\Phi} \mathrm{BU}.$$

It is the fibration induced from the universal fibration

$$(**) \quad U \rightarrow EU \rightarrow BU$$

by the map $1 - \psi^k : BU \rightarrow BU$. We consider the Serre spectral sequences of these fibrations. We know that

$$H^*(U; \mathbb{Z}) = \Lambda(y_1, y_2, \dots)$$

with $\deg y_i = 2i - 1$.

Since BU is simply connected we have

$$E_2^{r,s} = H^r(BU; \mathbb{Z}) \otimes H^s(U; \mathbb{Z}).$$

By [2] we know that the y_i 's are transgressive in $(**)$ with $d_{2i}y_i = \tilde{c}'_i \in E_{2i}^{2i,0}$. By naturality, the y_i are also transgressive in $(*)$ with

$$d_{2i}y_i = (1 - \psi^k)\tilde{c}'_i = (1 - k^i)\tilde{c}'_i \in E_{2i}^{2i,0}.$$

By induction on i , we see that in $(*)$ $E_{2i}^{*,*} = T_i \oplus L_i$ where T_i is a torsion group and L_i is \mathbb{Z} -free with basis the monomials $(\tilde{c}'_i)^{\alpha_i}(\tilde{c}'_{i+1})^{\alpha_{i+1}} \cdots y_i^{\beta_i} y_{i+1}^{\beta_{i+1}} \cdots$ with $\alpha_i \in \mathbb{N}$ and $\beta_i \in \{0, 1\}$. In particular, we have

$$E_{2i}^{2i,0} \cong \mathbb{Z}\tilde{c}'_i \oplus T \quad \text{and} \quad E_{\infty}^{2i,0} = E_{2i+1}^{2i,0} \cong \mathbb{Z}/s_i\tilde{c}'_i \oplus T,$$

where T is a torsion group and $s_i = |k^i - 1|$. Since $E_{\infty}^{2i,0} = \Phi^* H^{2i}(BU; \mathbb{Z})$, this proves Proposition 8.1. \square

Remark 8.2. This proof shows that \tilde{c}_i generates a direct summand of order s_i in $\text{Im } \Phi^*$ and that $\tilde{c}_{i-1}\tilde{c}_1$ is in $T \subset E_{2i}^{2i,0}$. Hence $\tilde{c}_i - \tilde{c}_{i-1}\tilde{c}_1 \in E_{\infty}^{2i,0} \subset \text{Im } \Phi^*$ has order s_i .

Let G_{∞} and SG_{∞} be the direct limits $\varinjlim G_n$ and $\varinjlim SG_n$ respectively. By performing the '+' construction on the classifying spaces we get the commutative diagram

$$\begin{array}{ccccc} BSG_n & \xrightarrow{f} & BSG_{\infty} & \longrightarrow & BSG_{\infty}^+ \\ \downarrow & & \downarrow j & & \downarrow \\ BG_n & \xrightarrow{g} & BG_{\infty} & \longrightarrow & BG_{\infty}^+ \sim JU(k) \end{array}$$

where the maps are the obvious maps.

Since $H^2(BSG_n; \mathbb{Z}/l) = 0$ for all $l \neq p$, we have $H^2(BSG_n; \mathbb{Z}[\frac{1}{p}]) = 0$. This proves that $\tilde{c}_1 = 0$ in $H^2(BSG_n; \mathbb{Z})$.

To prove the third part of Theorem 2.1, we observe that the map $BG_n \rightarrow JU(k)$

is a lifting of the map $BG_n \rightarrow BU$ obtained by the Brauer lifting of the canonical representation $BG_n \rightarrow BGL_n(\bar{\mathbb{F}}_p)$, where $\bar{\mathbb{F}}_p$ is an algebraic closure of \mathbb{F}_p .

Now if S is a l -Sylow subgroup of G_n or SG_n then l does not divide p and the Brauer character of the inclusion $S \rightarrow GL_n(\bar{\mathbb{F}}_p)$ is the character a complex representation of dimension n (cf. [6] and [8]). Hence $BS \rightarrow BG_n \rightarrow BU$ factorizes through $BU(n)$ and an argument of transfer gives the result.

The exact sequence $1 \rightarrow SG_\infty \rightarrow BG_\infty \xrightarrow{\det} G_1 \rightarrow 1$ splits and BG_∞^+ is an H-space (cf. [5]), so

$$BG_\infty^+ \text{ has the same homotopy type as } BSG_\infty^+ \times G_1 \quad (11)$$

(cf. also [1]). Thus in mod l cohomology, j^* is surjective and its kernel is the ideal generated by e_1 and c_1 .

By Theorems 1.1, 1.2 and 1.3, f^* and g^* are isomorphisms in mod l cohomology in degrees $\leq 2n$. Using the mapping cone of f and g , we see that f^* and g^* are injective in integral cohomology in these degrees away the p -torsion. But by [7] and [13], $H^*(BG_\infty; \mathbb{Z})$ has no p -torsion and then also $H^*(BSG_\infty; \mathbb{Z})$ by (11).

It remains to show that $j^* \tilde{c}_i$ has order $|k^i - 1|$. The monomorphism

$$h_n : G_{n-1} \rightarrow SG_n \subset G_n, \quad A \mapsto \begin{pmatrix} \det A^{-1} & 0 \\ 0 & A \end{pmatrix}$$

gives a map

$$h : BG_\infty \rightarrow BSG_\infty \xrightarrow{j} BG_\infty.$$

The following lemma and Remark 8.2 give the result. \square

Lemma 8.4. $h^* \tilde{c}_i = \tilde{c}_i - \tilde{c}_{i-1} \tilde{c}_1$.

Proof. The map $BG_n \rightarrow JU(k)$ is a lifting of the map $BG_n \rightarrow BU$ obtained by the Brauer lifting of the canonical representation $BG_n \rightarrow BGL_n(\bar{\mathbb{F}}_p)$, where $\bar{\mathbb{F}}_p$ is an algebraic closure of \mathbb{F}_p . Thus we can use the product formula to show that $h_n^* \tilde{c}_i = \tilde{c}_i - \tilde{c}_{i-1} \tilde{c}_1$ for $i < n$. The lemma follows from the injectivity of g^* . \square

9. Proof of Theorem 2.3

The groups $\tilde{H}^*(\bigotimes_{i \in I} C_i)$ and $\tilde{H}^*(BG; \mathbb{Z}[\frac{1}{p}])$ are of finite type. So by using the mapping cone of h and the universal coefficient theorem it is enough to show that h^* is an isomorphism modulo l for all primes l distinct from p . Then we consider only the i 's such that $i \equiv 0 \pmod{r}$. If \tilde{x} is an integral cohomology class, we denote by x its reduction modulo l .

It is easily seen that the homomorphism $H^*(C_i \otimes \mathbb{Z}/l) \rightarrow H^*(K_i; \mathbb{Z}/l)$, induced by the inclusion, is injective. Let $y_i \in H^{2i-1}(K_i; \mathbb{Z}/l)$ be the image of the generator $\bar{b}_i \in H^{2i-1}(C_i \otimes \mathbb{Z}/l) \cong \mathbb{Z}/l$. The monomials $x_i^\alpha y_i^\beta$, with $\alpha \in \mathbb{N}$ and $\beta \in \{0, 1\}$, form a basis of the image and we have $\beta y_i = \frac{s_i}{l} \tilde{x}_i$, where $\beta : H^{2i-1}(K_i; \mathbb{Z}/l) \rightarrow H^{2i}(K_i; \mathbb{Z})$ is the Bockstein homomorphism associated to the coefficient sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}/l \rightarrow 0$. But β is injective, so by the choice of \tilde{x}_i (cf. the proof of Lemma 2.2), y_i is homotopic to the map $K_i \rightarrow K(\mathbb{Z}/l, 2i-1)$ induced by the ring homomorphism $\mathbb{Z}/s_i \rightarrow \mathbb{Z}/l$. By the choice of γ_i we have then $\gamma_i^* y_i = e_i$, $\gamma_i^* \tilde{x}_i = \tilde{c}_i$ and $\gamma_i^* x_i = c_i$.

By identifying the cup products with the tensor products, we can regard $H^*(\otimes C_i \otimes \mathbb{Z}/l)$ as the subspace of $H^*(\prod K_i; \mathbb{Z}/l)$. The monomials

$$\omega = x_r^{\alpha_1} x_{2r}^{\alpha_2} \cdots x_{mr}^{\alpha_m} y_r^{\beta_1} y_{2r}^{\beta_2} \cdots y_{mr}^{\beta_m},$$

with $m = [n/r]$, form a basis of this subspace (if $G = \text{SG}_n$ and $r = 1$, the basis consists of the monomials $x_2^{\alpha_2} \cdots x_n^{\alpha_n} y_2^{\beta_2} \cdots y_n^{\beta_n}$). But the basis of $H^*(G; \mathbb{Z}/l)$ described in Section 1 consists of the monomials $g^* \omega$. This proves Theorem 2.3. \square

We prove Theorem 2.4 as above but we consider only the i 's such that $i \equiv 0 \pmod{2d}$. \square

Remark. For p odd we can obtain Theorem 2.4 as follows. The Adams operation ψ^q exists in symplectic K -theory for q odd and we have the fibration

$$J\text{Sp}(q) \xrightarrow{\Phi} B\text{Sp} \xrightarrow{1-\psi^q} B\text{Sp}.$$

We know that $H^*(B\text{Sp}; \mathbb{Z}) = \mathbb{Z}[\bar{c}'_2, \bar{c}'_4, \dots]$ where \bar{c}'_{2i} is the image of \tilde{c}'_{2i} by the canonical map $B\text{Sp} \rightarrow BU$. As before we show that $\Phi^* \bar{c}'_{2i}$ has order $q^{2i} - 1$ in $H^{4i}(J\text{Sp}(q); \mathbb{Z})$. By [7] there exists a commutative diagram

$$\begin{array}{ccc} B\text{Sp}_\infty & \xrightarrow{\chi_q} & J\text{Sp}(q) \\ \downarrow & & \downarrow \\ B\text{GL}_\infty & \xrightarrow{\chi_q} & JU(q) \end{array}$$

where $\chi_q^+ : B\text{Sp}_\infty^+ \rightarrow J\text{Sp}(q)$ is an homotopy equivalence (cf. also [5]). So Theorem 1.4 gives the mod l cohomology of $J\text{Sp}(q)$.

Since the diagram gives a "universal" class $\bar{e}_{2i} \in H^{4i-1}(J\text{Sp}(q); \mathbb{Z}/q^{2i} - 1)$, we can prove Theorem 2.4 as we proved Theorem 2.3. Finally, theorem 2.4 describes the integral cohomology of $J\text{Sp}(q)$ away the p -torsion, for p odd.

Acknowledgment

This paper was written during a stay at Cornell University. I wish to thank its members for their hospitality and especially Ken Brown. Some results of this paper are part of my thesis and I thank my adviser Ueli Suter for many stimulating discussions. I thank also the Swiss National Science Foundation for financial support.

References

- [1] D. Arlettaz, Chern-Klassen von ganzzahligen und rationalen Darstellungen diskreter Gruppen, *Math. Z.* 187 (1984) 49–60.
- [2] A. Borel, Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts, *Ann. of Math.* 57 (1953) 115–207.
- [3] K. Brown, *Cohomology of Groups* (Springer, Berlin, 1982).
- [4] H. Cartan et al., *Algebres d'Eilenberg–MacLane et homotopie*, Séminaire Henri Cartan, ENS, Paris, 1954/55.
- [5] Z. Fiedorowicz and S. Priddy, Homology of classical groups over finite fields and their associated infinite loop spaces, *Lecture Notes in Mathematics*, Vol. 674 (Springer, Berlin, 1978).
- [6] E.M. Friedlander, Unstable K -theories of the algebraic closure of a finite field, *Comment. Math. Helv.* 50 (1975) 145–154.
- [7] E.M. Friedlander, Computations of K -theories of finite fields, *Topology* 15 (1976) 87–109.
- [8] J.A. Green, The characters of the finite general linear groups, *Trans. Amer. Math. Soc.* 80 (1955) 402–447.
- [9] J. Huebschmann, The cohomology of $F\Psi^q$. The additive structure, *J. Pure Appl. Algebra* 45 (1987) 73–91.
- [10] D. Jeandupeux, Sur la cohomologie entière du groupe linéaire général et du groupe linéaire spécial sur un corps fini, *C.R. Acad. Sci. Paris Sér. I Math.* 308 (1989) 71–73.
- [11] D. Jeandupeux, *Cohomologie de produits semi-directs de groupes*, Thesis, Université de Neuchâtel, 1990.
- [12] D. Quillen, The Adams conjecture, *Topology* 10 (1971) 67–80.
- [13] D. Quillen, On the cohomology and K -theory of the general linear groups over a finite field, *Ann. of Math.* 96 (1972) 552–586.